

UNCLASSIFIED

Defense Technical Information Center
Compilation Part Notice

ADP013910

TITLE: Mathematical Models of Electromagnetic Wave Scattering by Two-Element Strip Grating with a Perpendicularly Magnetized Gyrotropic Medium

DISTRIBUTION: Approved for public release, distribution unlimited

This paper is part of the following report:

TITLE: 2002 International Conference on Mathematical Methods in Electromagnetic Theory [MMET 02]. Volume 2

To order the complete compilation report, use: ADA413455

The component part is provided here to allow users access to individually authored sections of proceedings, annals, symposia, etc. However, the component should be considered within the context of the overall compilation report and not as a stand-alone technical report.

The following component part numbers comprise the compilation report:

ADP013889 thru ADP013989

UNCLASSIFIED

MATHEMATICAL MODELS OF ELECTROMAGNETIC WAVE SCATTERING BY TWO-ELEMENT STRIP GRATING WITH A PERPENDICULARLY MAGNETIZED GYROTROPIC MEDIUM

Y.V. Gandel, V.V. Khoroshun
Kharkov National University
Pl. Svobody 4, Kharkov 61077, Ukraine
E-mail: gandel@ilt.kharkov.ua

ABSTRACT

Two mathematical models are proposed for analyzing a linearly polarized plane wave scattering by a strip grating placed on isotropic-gyrotropic media interface in the case of oblique incidence. The first mathematical model is based on reducing the original boundary value problem to the Riemann-Hilbert problem. The second model is based on reducing the same problem to a singular integral equation of the first kind with Cauchy kernel and its numerical solving by the discrete singularities method.

In this paper, the results obtained in [1-3] are generalized for the case of periodic structure consisting of two strips of different widths per period (two-element grating). This results in richer diffraction phenomena in comparison to simple grating because of additional control parameters. Moreover, unlike papers [1-3] here the case of oblique incidence of an H-polarized plane wave on a two-element grating is considered. The center of coordinate system is chosen in the middle of one of the strips.

The following set of dual series equations is mathematical model of a structure

$$\sum_n A_n \gamma_{n1} e^{in\varphi} = \gamma_{01}, \quad \theta_m < \varphi < \theta_m^{(1)}; \quad (1)$$

$$\sum_n A_n r_n e^{in\varphi} = -p_0, \quad \theta_m^{(1)} < \varphi < \theta_{m+1}^{(1)}, \quad (2)$$

where $\gamma_{n1} = \sqrt{k_0^2 n_1^2 - h_n^2}$, $n_1 = \sqrt{\epsilon_1 \mu_1}$, $h_n = k_0 n_1 \sin \zeta + \frac{2\pi}{l} n$,

$$r_n = 1 + \frac{\gamma_{n1}}{\epsilon_1 (R \gamma_{n2} - i L h_n)}, \quad \gamma_{n2} = \sqrt{k_0^2 \epsilon_{\perp} \mu_{\parallel} - h_n^2}, \quad \epsilon_{\perp} = \frac{\epsilon^2 - \epsilon_a^2}{\epsilon}$$

$$R = \epsilon_{\perp}^{-1}, \quad L = -\frac{\epsilon_a}{\epsilon^2 - \epsilon_a^2}, \quad k_0 = \frac{2\pi}{\lambda},$$

$$p_0 = 1 - \frac{\gamma_{01}}{\epsilon_1 (R \gamma_{02} - i L h_0)}, \quad \varphi = \frac{2\pi}{l} y, \quad \theta_m = 2\pi \frac{y_m}{l}, \quad (m=1,2),$$

l is the grating period, d is the slot width, λ is the wavelength, ζ is the incidence angle.

Denote $\tilde{A}_n = A_n r_n + p_0 \delta_{0n}$, and δ_{0n} for the Kronecker symbol.

Then initial set of dual series equations takes the form:

$$\sum_n \tilde{A}_n e^{in\varphi} = 0, \quad \theta_m < \varphi < \theta_m^{(1)}; \quad (3)$$

$$\sum_{n>0} \tilde{A}_n \frac{\eta_{n1}}{r_n^+} e^{m\varphi} + \sum_{n<0} |n| \tilde{A}_n \frac{\eta_{n1}}{r_n^-} e^{m\varphi} = (1 - \frac{\tilde{A}_0 - p_0}{r_0}) \kappa_1 \cos \zeta, \quad \theta_m^{(1)} < \varphi < \theta_{m+1}^{(1)}, \quad (4)$$

$$\sum_{n \neq 0} \tilde{A}_n e^{m\delta_m} = -\tilde{A}_0, \quad \theta_m^{(1)} < \delta_m < \theta_{m+1}^{(1)}, \quad (5)$$

The set of equations (3)-(5) can be reduced to non-homogeneous conjugation problem (Riemann-Hilbert's problem) with a complex-valued coefficient in the case of account of dissipative losses.

To calculate matrix elements of final matrix equation, it is necessary to introduce polynomials $Q_n(u_m, \rho)$ [3], where $\rho = \frac{\ln|G|}{2\pi}$, $u_m = \cos \theta_m$, ($m=1,2$).

In the case of multi-element gratings, an efficient numerical-analytical method for solving these dual series equations was suggested in [4]. The method consists in reducing them to a singular integral equation of the first kind with the Cauchy kernel on the set of segments, and its following solution by the method of discrete singularities [4,5]. Integral equation is of the following form

$$\frac{1}{\pi} \oint_L \frac{F(\xi)}{\xi - x} d\xi + \frac{1}{\pi} \int_L K(x, \xi) F(\xi) d\xi = f(x), \quad x \in L \quad (6)$$

where $L = \bigcup_{q=1}^m (a_q, b_q)$, $-\infty < a_1 < b_1 < \dots < a_m < b_m < +\infty$;

$f(x)$, $x \in \bar{L}$; $K(x, \xi)$, $x \in \bar{L}$, $\xi \in \bar{L}$ are known smooth functions, and function $F(\xi)$, $\xi \in L$ is sought in the functional class whose restriction on interval (a_q, b_q) :

$$F_q(\xi) = F(\xi), \quad a_q < \xi < b_q, \quad q = 1, \dots, m$$

can be represented in the form

$$F_q(\xi) = \frac{v_q(\xi)}{\sqrt{(\xi - a_q)(b_q - \xi)}}, \quad a_q < \xi < b_q,$$

where $v_q(\xi)$, $\xi \in [a_q, b_q]$ is a smooth function.

The sought function $F(\xi)$, $\xi \in L$ satisfies additional conditions, which in general case are of the following form:

$$\frac{1}{\pi} \int_L S_p(\xi) F(\xi) d\xi = C_p, \quad p = 1, \dots, m, \quad (7)$$

where $S_p(\xi)$, $\xi \in [a_p, b_p]$ is a known smooth function, and C_p is a known constant.

In conclusion we shall present the discrete mathematical model that is a set of linear algebraic equations for numerical solution of the integral equation (6) with additional condition (7).

Denote

$$t_i^n = \cos \frac{2i-1}{2n} \pi, \quad i = 1, \dots, n; \quad t_{0j}^n = \cos \frac{j}{n} \pi, \quad j = 1, \dots, n-1;$$

$$g_k(\tau) = \frac{b_k - a_k}{2}\tau + \frac{b_k + a_k}{2}; \quad \xi_{qi}^{n_q} = g_q(t_i^{n_q}), \quad i=1, \dots, n_q; q=1, \dots, m$$

$$\chi_{pj}^{n_p} = g_p(t_{0j}^{n_p}), \quad j=1, \dots, n_p-1; p=1, \dots, m$$

To calculate approximate values $\{v_{qn_q}(\xi)\}_{q=1}^m$ of the desired functions $v_q(\xi)$, $q=1, \dots, m$ in principal points $\{t_i^{n_q}\}_{i=1}^{n_q}$, we have a set of linear algebraic equations (where $R(x, \xi) = \frac{1}{\xi - x} + K(x, \xi)$)

$$\sum_{q=1}^m \sum_{i=1}^{n_q} R(\chi_{pj}^{n_p}, \xi_{qi}^{n_q}) v_{qn_q}(\xi_{qi}^{n_q}) \frac{1}{n_q} = f(\chi_{pj}^{n_p}), \quad j=1, \dots, n_p-1; p=1, \dots, m,$$

$$\sum_{i=1}^{n_p} S_p(\xi_{pi}^{n_p}) v_{pn_p}(\xi_{pi}^{n_p}) \frac{1}{n_p} = C_p, \quad (j=n_p), p=1, \dots, m$$

The values of the physical characteristic of scattered field,

$$H = \int_L H(\xi) F(\xi) d\xi = \sum_{q=1}^m \int_{a_q}^{b_q} H_q(\xi) v_q(\xi) \frac{d\xi}{\sqrt{(\xi - a_q)(b_q - \xi)}}.$$

are expressed in terms of the functions $v_q(\xi)$, $\xi \in [a_q, b_q]$, $q=1, \dots, m$,

where $H_q(\xi)$, $\xi \in [a_q, b_q]$ are known functions.

Approximate values of

$$H_{\bar{n}} = \sum_{q=1}^m \sum_{i=1}^{n_q} H_q(\xi_{qi}^{n_q}) v_{qn_q}(\xi_{qi}^{n_q}) \frac{1}{n_q}, \quad \bar{n} = (n_1, \dots, n_m)$$

are calculated in numerical experiments.

Obtained results can be applied in the design and elaboration of various devices containing periodic structures with ferrite substrates or in plasma.

REFERENCES

- [1] Y.V. Gandel, V.V. Khoroshun. Singular integral equations of the problem of electromagnetic wave diffraction by grating with a cross-magnetized ferrite, *Proc. Int. Symp. "Methods of Discrete Singularities in the Problems of Mathematical Physics"*, Kharkov, 1993, pp.123-124 (in Russian).
- [2] Y.V. Gandel, V.V. Khoroshun. The vortex lattice method in the electromagnetic wave diffraction on the method grating with gyrotropic layer, *Proc. Int. Conf. on Mathematical Methods in Electromagnetic Theory*, Kharkov, 2000, pp.578-580.
- [3] Y.V. Gandel, A.V. Litvinov, V.V. Khoroshun. Solution of the problem of electromagnetic wave diffraction by a strip grating with a cross-magnetized bigyrotropic medium, *Proc. Int. Symp. "Methods of Discrete Singularities in the Problems of Math. Phys."*, Orel, 2000, pp.128-132.
- [4] Y.V. Gandel. Method of discrete singularities in electromagnetic problems, *Cybernetics Problems*, M.: AN SSSR Publ., 1986, pp.166-183 (in Russian).
- [5] S.M. Belocerkovskiy, I.K. Lifanov. *Numerical Methods in Singular Integral Equations*, M.: Nauka, 1985 (in Russian).